arXiv:1610.03709v1 [math.AC] 12 Oct 2016

FINITE SUBGROUPS OF FIELDS

H.E.A. CAMPBELL, J. CHUAI, R.J. SHANK, AND D.L. WEHLAU

Abstract. Suppose $\mathbb{F}$ is a field of prime characteristic p and E is a finite subgroup of the additive group $(\mathbb{F}, +)$. Then E is an elementary abelian p-group. We consider two such subgroups, say E and E', to be equivalent if there is an $\alpha \in \mathbb{F}^* := \mathbb{F} \setminus \{0\}$ such that $E = \alpha E'$. In this paper we show that rational functions can be used to distinguish equivalence classes of subgroups and, for subgroups of prime rank or rank less than twelve, we give explicit finite sets of *separating invariants*.

1. Introduction

Suppose $\mathbb{F}$ is a field of prime characteristic p and E is a finite subgroup of the additive group $(\mathbb{F}, +)$. Then E is an elementary abelian p-group. We consider two such subgroups, say E and E', to be equivalent if there is an $\alpha \in \mathbb{F}^* := \mathbb{F} \setminus \{0\}$ such that $E = \alpha E'$. In this paper we show that rational functions can be used to distinguish equivalence classes of subgroups and, for subgroups of prime rank or rank less than twelve, we give explicit finite sets of *separating invariants*. Our motivation comes from Section 2 of [3], where the authors compute the rings of invariants for the two dimensional representations over $\mathbb{F}$ of elementary abelian p-groups. Every such representation is determined, up to group automorphism, by a finite subgroup of $(\mathbb{F}, +)$ and two such representations are equivalent if and only if the corresponding subgroups are in the same $\mathbb{F}^*$-orbit. The current paper is an attempt to understand the extent to which the ring of invariants determines the representation, at least for these types of representations.

2. Preliminaries

We start with a review of Section 2 of [3]. Define a group homomorphism $\rho : (\mathbb{F}, +) \to GL_2(\mathbb{F})$ by

$$\rho(c) := \begin{bmatrix} 1 & c \\ 0 & 1 \end{bmatrix}.$$

The restriction of ρ to a finite subgroup E gives a two dimensional representation of E. If E and E' are finite subgroups of $(\mathbb{F}, +)$, then $\rho(E)$ and $\rho(E')$ are conjugate if and only if there exists $\alpha \in \mathbb{F}^*$ with $E = \alpha E'$.

Date: October 13, 2016.

1991 *Mathematics Subject Classification.* 13A50.

This research was funded in part by NSERC and the University of New Brunswick.

There is a left action of $GL_2(\mathbb{F})$ on the polynomial ring $\mathbb{F}[x,y]$ defined by taking $x = [0\,1]$ and $y = [1\,0]$. For a finite subgroup $E \subset \mathbb{F}$, define

$$\mathbb{F}[x,y]^E := \{f \in \mathbb{F}[x,y] \mid f \cdot \rho(c) = f,\ \forall c \in E\}.$$

Clearly $x \in \mathbb{F}[x,y]^E$. A second invariant is given by

$$N_E(x,y) := \prod_{c\in E} y \cdot \rho(c) = \prod_{c\in E}(y + cx).$$

If the rank of E is r, then using the graded reverse lexicographic order with $y > x$, the leading term of $N_E(x,y)$ is y^{p^r}. Thus $\{x, N_E(x,y)\}$ is a homogeneous system of parameters for $\mathbb{F}[x,y]^E$ with $\deg(x)\deg(N_E(x,y)) = |E|$. Hence applying [4, 3.7.5] gives $\mathbb{F}[x,y]^E = \mathbb{F}[x, N_E(x,y)]$. Define

$$F_E(t) := \prod_{c\in E}(t - c) \in \mathbb{F}[t].$$

Thus $F_E(t)$ is the monic polynomial in t whose roots are the elements of E. Observe that $F_E(t) = N_E(1,t)$ and $N_E(x,y) = x^{|E|}F_E(y/x)$. Furthermore, for $\alpha \in \mathbb{F}^*$, we have $F_{\alpha E}(t) = \alpha^{|E|}F_E(t/\alpha)$.

If $\{x_1, \dots, x_r\}$ is a basis for $(\mathbb{F}_p{}^r)^*$ then

$$D_r(t) := \prod_{u\in(\mathbb{F}_p{}^r)^*} (t - u) \in \mathbb{F}_p[x_1, \dots, x_r]^{GL_r(\mathbb{F}_p)}[t].$$

It is well-known that $D_r(t) = t^{p^r} + \sum_{i=0}^{r-1} d_{r-i,r}t^{p^i}$, where the $d_{j,r}$ are the Dickson invariants, and

$$\mathbb{F}_p[x_1, \dots, x_r]^{GL_r(\mathbb{F}_p)} = \mathbb{F}_p[d_{1,r}, \dots, d_{r,r}]$$

(see, for example, [6]). Using the inclusion $\mathbb{F}_p \subset \mathbb{F}$, we can interpret elements of $\mathbb{F}_p[x_1, \dots, x_r]$ as polynomial functions on $\mathbb{F}^r$. If $\{c_1, \dots, c_r\} \subset \mathbb{F}$ is an $\mathbb{F}_p$-basis for E, then $d_{i,r}(c_1, \dots, c_r) \in \mathbb{F}$. Since $d_{i,r} \in \mathbb{F}_p[x_1, \dots, x_r]^{GL_r(\mathbb{F}_p)}$, the result is independent of the choice of basis for E and so it makes sense to define $d_{i,r}(E) := d_{i,r}(c_1, \dots, c_r)$. Since evaluating the coefficients of $D_r(t)$ at a basis for E gives $F_E(t)$, we have the following.

Proposition 2.1. $N_E(x,y) = y^{p^r} + \sum_{i=0}^{r-1} d_{i-r,r}(E)y^{p^i}x^{p^r-p^i}$.

Note that, since $d_{r,r}(b_1, \dots, b_r)$ is the product of the $\mathbb{F}_p$-linear combinations of $\{b_1, \dots, b_r\}$, we have $\operatorname{rank}(\operatorname{Span}_{\mathbb{F}_p}\{b_1, \dots, b_r\}) = r$ if and only if $d_{r,r}(b_1, \dots, b_r) \neq 0$.

Our goal is to use rational functions to distinguish equivalence classes of finite subgroups. A point $(c_1, \dots, c_r) \in \mathbb{F}^r$ determines a finite subgroup $E = \operatorname{Span}_{\mathbb{F}_p}\{c_1, \dots, c_r\}$. The rank of E is r if and only if $d_{r,r}(E) \neq 0$. Two points determine the same subgroup if they lie is a common $GL_r(\mathbb{F}_p)$-orbit. Thus equivalence classes of rank r subgroups correspond to points in the orbit space $GL_r(\mathbb{F}_p)\backslash\mathbb{F}^r/\mathbb{F}^*$ with $d_{r,r}(c_1, \dots, c_r) \neq 0$. Rather than working with the Zariski open set $d_{r,r} \neq 0$, it is convenient to introduce a formal inverse for $d_{r,r}$, say t, and study the resulting Zariski closed set $d_{r,r}t = 1$. Since $d_{i,r}$ is homogeneous of degree $p^r - p^{r-i}$, for $\alpha \in \mathbb{F}^*$ we have $d_{i,r}(\alpha E) = \alpha^{p^r-p^{r-i}}d_{i,r}(E)$. Thus $\mathbb{F}^*$

acts on $d_{i,r}$ with weight $p^r - p^{r-i}$. To ensure that the equation $d_{r,r}t = 1$ is isobaric with weight zero, we assign weight $1 - p^r$ to t. This defines an action of $\mathbb{F}^*$ on $\mathbb{F}[d_{1,r}, \ldots, d_{r,r}, t]$.

Proposition 2.2. *Suppose $\mathbb{F}$ is algebraically closed. Then the polynomial functions $\mathbb{F}[d_{1,r}, \ldots, d_{r,r}, t]^{\mathbb{F}^*}$ separate the orbits satisfying $d_{r,r}t = 1$. Furthermore, $\mathbb{F}[d_{1,r}, \ldots, d_{r,r}, t]^{\mathbb{F}^*}$ is generated by $d_{r,r}t$ and a finite set of monomials in $\mathbb{F}[d_{1,r}, \ldots, d_{r-1,r}, t]$. Thus equivalence classes of rank r subgroups are distinguished by a finite set of rational functions given by monomials in $\mathbb{F}[d_{1,r}, \ldots, d_{r-1,r}, d_{r,r}^{-1}]$.*

Proof. The multiplicative group $\mathbb{F}^*$ is linearly reductive (see, for example [4, Theorem 2.2.19]). Therefore the ring of invariants is finitely generated and the invariants separate closed orbits. For an $\mathbb{F}^*$-action, if an orbit is not closed then the closure includes one additional point. This point must be a closed orbit and hence a fixed point. For the given action, the only fixed point is the the zero vector. Thus the invariants separate the orbits satisfying $d_{r,r}t = 1$. Since the action is diagonal, the invariants are generated by a set of monomials. Note that the weight of $d_{i,r}$ for $i = 1, \ldots, r$ is positive. Thus every monomial of positive degree and weight zero involves a positive power of t. □

Definition 2.3. For a finite subgroup of rank r, the following $\mathbb{F}^*$-invariant rational functions are well-defined:

$$v_1 := \frac{d_{1,r}^{(p^r-1)/(p-1)}}{d_{r,r}^{p^{r-1}}} \text{ and } v_{1j} := \frac{d_{j,r}d_{1,r}^{p(p^{r-j}-1)/(p-1)}}{d_{r,r}^{p^{r-j}}}$$

for $j \in \{2, \ldots, r-1\}$.

Groups of Rank 2. Subgroups of rank 2 provide an instructive special case. The group $\mathbb{F}^*$ acts on $d_{1,2}$ with weight $p^2 - p$, $d_{2,2}$ with weight $p^2 - 1$ and t with with weight $1 - p^2$. It is clear that

$$\mathbb{F}[d_{1,2}, d_{2,2}, t]^{\mathbb{F}^*} = \mathbb{F}[d_{1,2}^{p+1}t^p, d_{2,2}t].$$

The groups of rank 2 correspond to orbits with $d_{2,2}t = 1$ and the orbits of rank 2 subgroups are separated by the rational function $v_1 = d_{1,2}^{p+1}/d_{2,2}^p$. Note that $v_1 = v_1(x, y)$ is a rational function of x, y and $v_1(a, b)$ is defined whenever $\{a, b\}$ is linearly independent over $\mathbb{F}_p$. If V is a rank 2 subgroup of $\mathbb{F}$ then we will write $v_1(V) := d_{1,2}^{p+1}(V)/d_{2,2}^p(V)$.

In 1911, Dickson [5] showed that $d_{1,2}(x, y) = -\prod_{(a,b)\in\Omega} ax^2+baxy+y^2$ where the product is over all $a, b \in \mathbb{F}_p$ such that z^2+bz+a is an irreducible quadratic in $\mathbb{F}_p[z]$. Thus $d_{1,2}(1, \gamma) = 0$ if and only if γ satisfies an irreducible quadratic over $\mathbb{F}_p$. In this case, $\mathrm{Span}_{\mathbb{F}_p}\{1, \gamma\} = \mathbb{F}_{p^2}$. More generally if $\alpha \neq 0$ then $d_{1,2}(\alpha, \beta) = \alpha^{p^2-p}d_{1,2}(1, \beta/\alpha) = 0$ if and only if β/α satisfies an irreducible quadratic over $\mathbb{F}_p$. If $\{\alpha, \beta\}$ is any $\mathbb{F}_p$-basis of V then $\{1, \beta/\alpha\}$ is a basis of $\alpha^{-1} \cdot V$. Thus $v_1(V) = 0$ if and only if $v_1(\alpha^{-1} \cdot V) = 0$ if and only if $d_{1,2}(1, \beta/\alpha) = 0$ if and only if $\alpha^{-1} \cdot V = \mathbb{F}_{p^2}$. In other words, the $\mathbb{F}^*$-orbit

determined by $v_1(V) = 0$ consists of the dilations of $\mathbb{F}_{p^2} \subset \mathbb{F}$. This example suggests that some finite subgroups possess additional structure.

3. The Structure of a Subgroup

For a finite subgroup $E \subset (\mathbb{F}, +)$, define the stabiliser

$$\mathrm{Stab}_{\mathbb{F}^*}(E) := \{\alpha \in \mathbb{F}^* \mid \alpha E = E\}.$$

Proposition 3.1. *Suppose E a is a finite subgroup of $\mathbb{F}$ of rank r. Then $\mathrm{Stab}_{\mathbb{F}^*}(E) = \mathbb{F}_q^*$ for some finite subfield $\mathbb{F}_q \subseteq \mathbb{F}$ with $q \leq p^r$. Furthermore, E is a vector space over $\mathbb{F}_{p^s}$ if and only if $\mathbb{F}_{p^s} \subseteq \mathbb{F}_q$.*

Proof. Clearly $\mathrm{Stab}_{\mathbb{F}^*}(E)$ is closed under multiplication and contains 1. Define $L := \mathrm{Stab}_{\mathbb{F}^*}(E) \cup \{0\}$. For $\alpha_1, \alpha_2 \in L$ and $\gamma \in E$,

$$(\alpha_1 + \alpha_2)\gamma = \alpha_1\gamma + \alpha_2\gamma \in E.$$

Thus L is a subfield of $\mathbb{F}$. Choose $\gamma \in E \setminus \{0\}$. Then $1 \in \gamma^{-1}E$ and $\mathrm{Stab}_{\mathbb{F}^*}(\gamma^{-1}E) = \mathrm{Stab}_{\mathbb{F}^*}(E)$. Therefore $L \subseteq \gamma^{-1}E$, $q := |L| \leq |E| = p^r$ and E is a vector space over $L = \mathbb{F}_q$. Furthermore, if E is a vector space over $\mathbb{F}_{p^s}$, then $\mathbb{F}_{p^s}^* \subseteq \mathrm{Stab}_{\mathbb{F}^*}(E) = \mathbb{F}_q^*$. □

Theorem 3.2. *Suppose $q = p^s$ and $\mathbb{F}_q \subset \mathbb{F}$. Further suppose that E is a finite subgroup of $\mathbb{F}$ of rank $r = ms$ for some integer m. Then E is a vector space over $\mathbb{F}_q$ if and only if $d_{i,r}(E) = 0$ for all i not congruent to zero modulo s.*

Proof. Denote $\hom_{\mathbb{F}_q}({\mathbb{F}_q}^m, \mathbb{F}_q)$ by $({\mathbb{F}_q}^m)^*$ and define

$$D_{m,q}(t) := \prod_{u \in ({\mathbb{F}_q}^m)^*} (t - u) \in \mathbb{F}_q[y_1, \ldots, y_m]^{GL_m(\mathbb{F}_q)}[t].$$

The coefficients of $D_{m,q}(t)$ are the Dickson invariants over the field $\mathbb{F}_q$. Thus $D_{m,q}(t) = t^{q^m} + \sum_{j=0}^{m-1} d_{m-j,m}^{(q)} t^{q^j}$.

Suppose E is a vector space over $\mathbb{F}_q$ with basis $\{b_1, \ldots, b_m\}$. Then evaluating the coefficients of $D_{m,q}(t)$ at $(b_1, \ldots, b_m)$ gives a monic polynomial in $\mathbb{F}[t]$ whose roots are the elements of E, in other words, we recover $F_E(t)$. We can also construct $F_E(t)$ by evaluating the coefficients of $D_r(t)$. By comparing these constructions, we see that $d_{r-\ell,r}(E) = 0$ unless $\ell = js$. Hence $d_{i,r}(E) = 0$ unless $i = js - ms = s(m - j)$.

Suppose $d_{i,r}(E) = 0$ for all i not congruent to zero modulo s. We will show that $\mathbb{F}_{p^s} \subseteq \mathrm{Stab}_{\mathbb{F}^*}(E)$ and, therefore, using Proposition 3.1, E is a subspace over $\mathbb{F}_q$. The polynomial $d_{i,r}$ is homogeneous of degree $p^r - p^{r-i} = p^{r-i}(p^i - 1)$. If $i = sk$ then $p^s - 1$ divides $p^{ks} - 1 = p^i - 1$. For $\alpha \in \mathbb{F}_{p^s}^*$, we have $\alpha^{p^s-1} = 1$ and $d_{ks,r}(\alpha E) = d_{ks,r}(E)$. Furthermore, if $d_{i,r}(E) = 0$ then $d_{i,r}(\alpha E) = 0$. Thus, if $d_{i,r}(E) = 0$ for all i not divisible by s, we have $d_{i,r}(\alpha E) = d_{i,r}(E)$ for all i and all $\alpha \in \mathbb{F}_{p^s}^*$. The Dickson invariants separate $GL_r(\mathbb{F}_p)$-orbits. Therefore, if $d_{i,r}(\alpha E) = d_{i,r}(E)$ for all i, then E and αE are in the same $GL_r(\mathbb{F}_p)$-orbit,

i.e., if $\{u_1, ..., u_r\}$ is a basis for E, then $\{\alpha u_1, ..., \alpha u_r\}$ is also a basis for E. Thus if $\alpha \in \mathbb{F}_{p^s}$ then α stabilises E. □

Among the subgroups of $(\mathbb{F}, +)$, we distinguish the dilations of finite fields: $V_{\alpha,r} := \alpha \cdot \mathbb{F}_{p^r}$ where $\alpha \in \mathbb{F}^*$ and r is a positive integer. Any finite subgroup $V \subseteq (\mathbb{F}, +)$ decomposes as a direct sum

$$V = \bigoplus_{j=1}^{s} V_{\alpha_j,\lambda_j}$$

where $\alpha_j \in \mathbb{F}^*$ and $\lambda_j \geq 1$. For example, if $\{\alpha_1, \ldots, \alpha_s\}$ is an $\mathbb{F}_p$-basis for V then $V = \oplus_{j=1}^{s} V_{\alpha_j,1}$. The decomposition is not necessarily unique. If $r = m \cdot \ell$, then $\mathbb{F}_{p^r}$ is an $\mathbb{F}_{p^\ell}$ vector space of dimension m. If $\{e_1, \ldots, e_m\}$ is an $\mathbb{F}_{p^\ell}$-basis for $\mathbb{F}_{p^r}$, then $V_{\alpha,r} = \oplus_{i=1}^{m} V_{\alpha e_i,\ell}$. For each decomposition we define the sequence $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_s)$. The sequence is a partition of the integer $\dim_{\mathbb{F}_p}(V)$. If we order the sequences lexicographically from the left, then for each V there is a unique maximal sequence with $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_s$. We will denote this sequence by $\lambda(V)$ and call the sequence the *partition* of V, i.e., the partition of V is the largest, with respect to lex order, of the sequences which correspond to decompositions of V. If V has rank 2, then $\lambda(V)$ is either (2) or $(1, 1)$ and $\lambda(V) = 2$ if and only if $(d_{1,2}^{p+1}/d_{2,2}^{p})(V) = 0$.

Define a function $\omega_s : \mathbb{F} \to \mathbb{F}$ by $\omega_s(t) = t^{p^s} - t$. Then ω_s is $\mathbb{F}_p$-linear and the kernel of ω_s is $\mathbb{F}_{p^s}$. Furthermore, $\omega_s(a) = \omega_s(b)$ if and only if $a - b \in \mathbb{F}_{p^s}$. Therefore, if V has rank r and $V \cap \mathbb{F}_{p^s} = \{0\}$ then $\omega_s(V)$ is a finite subgroup of rank r. In the following, we use the convention that $d_{i,r}(V) = 0$ if $i < 0$ or $i > \text{rank}(V)$.

Theorem 3.3. *If V is a finite subgroup of rank r and $V = \mathbf{F}_{p^s} \oplus W$, where W is a finite subgroup of rank $\ell = r - s$, then*

$$d_{i,r}(V) = d_{i,\ell}\left(\omega_s\left(W\right)\right) - d_{i-s,\ell}\left(\omega_s\left(W\right)\right).$$

Proof. Recall that

$$N_V(x, y) = \sum_{i=0}^{r} d_{r-i,r}(V) y^{p^i} x^{p^r - p^i}.$$

Furthermore, since the elements of $\mathbb{F}_{p^s}$ are the roots of $t^{p^s} - t$, we have $N_{\mathbb{F}_{p^s}}(x, y) = y^{p^s} - y x^{p^s - 1}$. Thus $d_{s,s}(\mathbb{F}_{p^s}) = -1$ and $d_{i,s}(\mathbb{F}_{p^s}) = 0$ for $0 < i < s$.

Using the decomposition of V, we get

$$\begin{aligned}
N_V(x,y) &= \prod_{v\in V}(y+vx) = \prod_{a\in\mathbb{F}_{p^s},b\in W}(y+(a+b)x)\\
&= \prod_{b\in W}\left(\prod_{a\in\mathbb{F}_{p^s}}((y+bx)+ax)\right)\\
&= \prod_{b\in W}N_{\mathbb{F}_{p^s}}(x,y+bx) = \prod_{b\in W}\left((y+bx)^{p^s}-(y+bx)x^{p^s-1}\right)\\
&= \left(x^{p^s-1}\right)^{p^{r-s}}\prod_{b\in W}\left(\left(\frac{y^{p^s}}{x^{p^s-1}}-y\right)+(b^{p^s}-b)x\right)\\
&= \left(x^{p^s-1}\right)^{p^{r-s}}N_{\omega_s(W)}\left(x,\frac{y^{p^s}}{x^{p^s-1}}-y\right).
\end{aligned}$$

Since $N_E(x,y)$ is an $\mathbb{F}_p$-linear function of y,

$$\begin{aligned}
N_V(x,y) &= \left(x^{p^s-1}\right)^{p^{r-s}}\left(N_{\omega_s(W)}\left(x,\frac{y^{p^s}}{x^{p^s-1}}\right)-N_{\omega_s(W)}(x,y)\right)\\
&= \left(\sum_{i=0}^{r-s}d_{r-s-i,\ell}\left(\omega_s\left(W\right)\right)y^{p^{s+i}}x^{p^r-p^{s+i}}\right)-\left(\sum_{i=0}^{r-s}d_{r-s-i,\ell}\left(\omega_s\left(W\right)\right)y^{p^i}x^{p^r-p^i}\right)\\
&= \left(\sum_{j=s}^{r}d_{r-j,\ell}\left(\omega_s\left(W\right)\right)y^{p^j}x^{p^r-p^j}\right)-\left(\sum_{i=0}^{r-s}d_{r-s-i,\ell}\left(\omega_s\left(W\right)\right)y^{p^i}x^{p^r-p^i}\right)\\
&= \sum_{k=0}^{r}\left(d_{r-k,\ell}\left(\omega_s\left(W\right)\right)-d_{r-s-k,\ell}\left(\omega_s\left(W\right)\right)\right)y^{p^k}x^{p^r-p^k}.
\end{aligned}$$

Thus $d_{i,r}(V)=d_{i,\ell}\left(\omega_s\left(W\right)\right)-d_{i-s,\ell}\left(\omega_s\left(W\right)\right)$. □

Theorem 3.4. *Suppose V is a finite subgroup of rank $s+1$ with $s>1$. Then the partition of V is $(s,1)$ if and only if $d_{i,s+1}(V)=0$ for $1<i<s$ and*

$$\left(\frac{d_{1,s+1}^p d_{s,s+1}}{d_{s+1,s+1}^p}\right)(V)=1.$$

Proof. Suppose V has partition $(s,1)$. Then $\alpha V=\mathbb{F}_{p^s}\oplus\beta F_p$ for some $\alpha\in\mathbb{F}^*$ and $\beta\in\mathbb{F}\setminus\mathbb{F}_{p^s}$. As in section 2, denote $v_{1s}=(d_{1,s+1}^p d_{s,s+1})/(d_{s+1,s+1}^p)$. Since v_{1s} is $\mathbb{F}^*$-invariant, $v_{1s}(V)=v_{1s}(\mathbb{F}_{p^s}\oplus\beta F_p)$. Furthermore, $d_{i,s+1}(V)=0$ if and only if $d_{i,s+1}(\alpha V)=0$. Therefore, using Theorem 3.3, $d_{i,s+1}(V)=0$ for $0<i<s$, $d_{1,s+1}(V)=-d_{s+1,s+1}(V)$ and $d_{s,s+1}(V)=-1$, giving $v_{1s}(V)=1$ as required.

Define $f:=d_{s+1,s+1}^p-d_{1,s+1}^p d_{s,s+1}$. We will show that if $f(V)=0$ and $d_{i,s+1}(V)=0$ for $0<i<s$, then V contains a subgroup equivalent to $\mathbb{F}_{p^s}$.

Suppose V has basis $\{c_1,\ldots,c_{s+1}\}$. Using Theorem 3.2, if $d_{i,s}(c_1,\ldots,c_s)=0$ for $0<i<s$, then $\mathrm{Span}_{\mathbb{F}_p}\{c_1,\ldots,c_s\}$ is equivalent to $\mathbb{F}_{p^s}$. Thus, to show

that the partition of V is $(s,1)$, it is sufficient to find a $g \in GL_{s+1}(\mathbb{F}_p)$ such that $d_{i,s}g(c_1,\ldots,c_{s+1}) = 0$ for $0 < i < s$. (Since $d_{i,s} \in \mathbb{F}_p[x_1,\ldots,x_s] \subset \mathbb{F}[x_1,\ldots,x_{s+1}]$, we can evaluate $d_{i,s}$ on an element of $\mathbb{F}^{s+1}$.)

Recall that $d_{i,s+1} = d_{i,s}^p - d_{i-1,s}D_s^{p-1}(x_{s+1})$ (see Proposition 1.3(b) of [6]). In particular $d_{1,s+1} = d_{1,s}^p - D_s^{p-1}(x_{s+1})$ and $d_{s+1,s+1} = -d_{s,s}D_s^{p-1}(x_{s+1})$. Suppose there is a basis $\{b_1,\ldots,b_{s+1}\}$ for V with $d_{1,s}(b_1,\ldots,b_{s+1}) = 0$. Then, since $d_{2,s}^p = d_{2,s+1} + d_{1,s}D_s^{p-1}(x_{s+1})$, we have $d_{2,s}(b_1,\ldots,b_{s+1}) = 0$. Similarly, $d_{i,s}(b_1,\ldots,b_{s+1}) = 0$ for $i = 3,\ldots,s-1$. Thus it is sufficient to find a basis for which $d_{1,s}(b_1,\ldots,b_{s+1}) = 0$.

Let Π_s denote the $GL_{s+1}(\mathbb{F}_p)$-orbit product of $d_{1,s}$. The degree of Π_s is $\deg(d_{1,s})(p^{s+1}-1)/(p-1) = p^{s-1}(p^{s+1}-1)$ (look at the stabiliser of the hyperplane $x_{s+1} = 0$). Define

$$H := f^{p^{s-2}} + \sum_{j=1}^{s-2}(-1)^{j+1}d_{s-j,s+1}^{p^{s-2-j}}d_{1,s+1}^{p^{(s-1-j)}(p^{j+1}-1)/(p-1)}.$$

Note that H is non-zero and homogeneous of degree $p^{s-1}(p^{s+1}-1)$. We will show that $d_{1,s}$ divides H and, therefore, H is non-zero scalar multiple of Π_s. Since $H(V) = 0$, we then conclude that $\Pi_s(V) = 0$ and, therefore, $d_{1,s}(gV) = 0$ for some g. Hence the partition of V is $(s,1)$.

We work modulo the ideal $\mathfrak{n} := \langle d_{1,s}\rangle$. We will use induction on i to prove that

$$f^{p^i} + \sum_{j=1}^{i}(-1)^{j+1}d_{s-j,s+1}^{p^{i-j}}d_{1,s+1}^{p^{(i-j+1)}(p^{j+1}-1)/(p-1)} \equiv_{\mathfrak{n}} (-1)^{i+1}d_{s-1-i,s}d_{1,s+1}^{(p^{i+2}-1)/(p-1)}$$

for $i \in \{0,\ldots,s-2\}$. For $i = s-2$, the left side of this equivalence is H and the right is congruent to zero.

For convenience, we use h to denote $D_s^{p-1}(x_{s+1})$. Thus

$$d_{i,s+1} = d_{i,s}^p - d_{i-1,s}h \in \mathbb{F}_p[d_{1,s},\ldots,d_{i,s},h]$$

and $d_{1,s+1} \equiv_{\mathfrak{n}} -h$. Using this equivalence, $d_{s+1,s+1} = -d_{s,s}h \equiv_{\mathfrak{n}} d_{s,s}d_{1,s+1}$ and $d_{s,s+1} \equiv_{\mathfrak{n}} d_{s,s}^p + d_{s-1,s}d_{1,s+1}$. Thus

$$f = d_{s+1,s+1}^p - d_{1,s+1}^p d_{s,s+1} \equiv_{\mathfrak{n}} d_{s,s}^p d_{1,s+1}^p - d_{1,s+1}^p\left(d_{s,s}^p + d_{s-1,s}d_{1,s+1}\right) = -d_{s-1,s}d_{1,d+1}^{p+1}$$

which is the $i = 0$ case.

Assume the result is true for i and take the p^{th} power:

$$f^{p^{i+1}} + \sum_{j=1}^{i}(-1)^{j+1}d_{s-j,s+1}^{p^{i-j+1}}d_{1,s+1}^{p^{(i-j+2)}(p^{j+1}-1)/(p-1)} \equiv_{\mathfrak{n}} (-1)^{i+1}d_{s-1-i,s}^{p}d_{1,s+1}^{p(p^{i+2}-1)/(p-1)}.$$

However, $d^p_{s-i-1,s} \equiv_{\mathfrak{n}} d_{s-1-i,s+1} - d_{1,s+1}d_{s-i-2,s}$. Substituting this into the right side of the equivalence and bringing one of the resulting terms to the left gives

$$\begin{aligned} f^{p^{i+1}} \quad &+ \quad \sum_{j=1}^{i+1}(-1)^{j+1} d^{p^{i+1-j}}_{s-j,s+1} d^{p^{((i+1)-j+1)}(p^{j+1}-1)/(p-1)}_{1,s+1} \\ &\equiv_{\mathfrak{n}} \quad (-1)^{(i+1)+1} d_{s-1-(i-1),s} d^{(p^{(i+1)+2}-1)/(p-1)}_{1,s+1}, \end{aligned}$$

as required. □

Theorem 3.5. *Suppose $\mathbb{F}$ is algebraically closed and V is a finite subgroup of rank $r = s-1$. Then V is a subgroup of a dilation of $\mathbb{F}_{p^s}$ if and only if $d_{1,r}(V) \neq 0$, $v^p_{12}(V) = v_1(V)$ and $v^p_{1i}(V) = v_{1(i-1)}(V)$ for $i \in \{3, \dots, r-1\}$.*

Proof. As in the proof of Theorem 3.4, we use the equations

$$d_{i,s} = d^p_{i,r} - d_{i-1,r} D^{p-1}_r(x_s).$$

Choose a basis $\{c_1, \dots, c_r\}$ for V. Note that $D_r(t) \in \mathbb{F}[x_1, \dots, x_r][t]$ and let $\widetilde{D_r}(t)$ denote the polynomial in $\mathbb{F}[t]$ constructed by setting x_i to c_i in $D_r(t)$. Since $\mathbb{F}$ is algebraically closed, $\mathbb{F}$ contains a root of $\widetilde{D_r}(t)^{p-1} - d^p_{1,r}(V)$, say c_s. Take E to be the $\mathbb{F}_p$-span of $\{c_1, \dots, c_s\}$. Thus $\widetilde{D_r}(c_s)^{p-1} = d^p_{1,r}(V)$ and $d_{i,s}(E) = \left(d^p_{i,r} - d_{i-1,r}d^p_{1,r}\right)(V)$. In particular, $d_{s,s}(E) = -d_{r,r}(V)d_{1,r}(V)$. Hence E has rank s if and only if $d_{1,r}(V) \neq 0$.

Using Theorem 3.2, E is a dilation of $\mathbb{F}_{p^s}$ if and only if $d_{1,r}(V) \neq 0$ and $0 = d_{i,s}(E) = \left(d^p_{i,r} - d_{i-1,r}d^p_{1,r}\right)(V)$ for $i \in \{1, \dots, s-1\}$. For $i > 2$, if we multiply $d^p_{i,r} - d_{i-1,r}d^p_{1,r}$ by $d^{p^2(p^{r-i}-1)/(p-1)}_{1,r} / d^{p-i+1}_{r,r}$, we get the $\mathbb{F}^*$-invariant form $v^p_{1i} - v_{1(i-1)}$. For $i = 2$, the invariant form is $v^p_{12} - v_1$. Thus, if $d_{1,r}(V) \neq 0$, $(v^p_{12} - v_1)(V) = 0$ and $\left(v^p_{1i} - v_{1(i-1)}\right)(V) = 0$ for $i \in \{3, \dots, r-1\}$, V is contained in a dilation of $\mathbb{F}_{p^s}$. To prove the converse, observe that if $V \subset W$ and W is a dilation of $\mathbb{F}_{p^s}$ then the equations are easily seen to be satisfied. □

4. Separating Invariants

It follows from Proposition 2.2 that, for $\mathbb{F}$ algebraically closed, the orbits of finite subgroups of rank r are separated by the rational functions

$$\mathbb{F}[d_{1,r}, \dots, d_{r-1,r}, d^{-1}_{r,r}]^{\mathbb{F}^*}.$$

In the following, whenever we consider separating invariants, we assume $\mathbb{F}$ is algebraically closed. The ring $\mathbb{F}[d_{1,r}, \dots, d_{r-1,r}, d^{-1}_{r,r}]^{\mathbb{F}^*}$ is spanned by the monomials $d^{a_1}_{1,r} \cdots d^{a_{r-1}}_{r-1,r} d^{-a_r}_{r,r}$ satisfying the equation

$$a_1 p^{r-1}(p-1) + a_2 p^{r-2}(p^2-1) + \cdots + a_{r-1}p(p^{r-1}-1) = a_r(p^r - 1). \tag{4.1}$$

The non-negative integer solutions to Equation 4.1 form a monoid. This monoid inherits a partial order from the integer lattice $\mathbb{Z}^r$. We will refer to a non-zero element of the monoid as *primitive* if it can not be written a sum of two smaller elements. We will refer to a monomial with a primitive

exponent sequence as a primitive invariant. It is clear that the set of primitive elements in the monoid give the exponents of a generating set for the ring of invariants. We will often abuse notation by using the same symbol to denote the sequence $(a_1, \dots, a_r)$ and the monomial $d_{1,r}^{a_1} \cdots d_{r-1,r}^{a_{r-1}} d_{r,r}^{-a_r}$. For every $i < r$, let ℓ denote the greatest common divisor of $p^i - 1$ and $p^r - 1$. Then

$$v_i := d_{i,r}^a d_{r,r}^{-b},$$

with $a = (p^r - 1)/\ell$ and $b = p^{r-i}(p^i - 1)/\ell$, is the unique primitive invariant with support $\{d_{i,r}, d_{r,r}^{-1}\}$. Note that this definition of v_i is consistent with the definition of v_1 given in Definition 2.3.

If $(a_1, \dots, a_r)$ is solution to Equation 4.1, we will refer to $a_1 + a_2 + \cdots + a_{r-1} - a_r$ as the *height* of the solution. This is just the degree of the monomial $d_{1,r}^{a_1} \cdots d_{r-1,r}^{a_{r-1}} d_{r,r}^{-a_r}$ if we assign degree one to each $d_{i,r}$. Thus a primitive solution has height at least one and any non-negative solution of height one is primitive.

Lemma 4.1. *If the greatest common divisor of r and i divides j, then there exists a primitive invariant*

$$v_{ij} := d_{j,r} d_{i,r}^a d_{r,r}^{-b}.$$

Proof. To find invariants with support $\{d_{i,r}, d_{j,r}, d_{r,r}^{-1}\}$, we consider the equation

$$a_j p^{r-j}(p^j - 1) + a_i p^{r-i}(p^i - 1) = a_r(p^r - 1).$$

Suppose m is the greatest common divisor of i and r. Then $p^m - 1$ is the greatest common divisor of $p^r - 1$ and $p^{r-i}(p^i - 1)$. Thus there exist positive integers $\bar{a}$ and $\bar{b}$ with $p^m - 1 = \bar{b}(p^r - 1) - \bar{a}p^{r-i}(p^i - 1)$.

Since m divides j, $p^m - 1$ divides $p^j - 1$. Thus $c := p^{r-j}(p^j - 1)/(p^m - 1)$ is a positive integer. Define $a := c\bar{a}$ and $b := c\bar{b}$. Then $p^{r-j}(p^j - 1) = b(p^r - 1) - ap^{r-i}(p^i - 1)$. Hence $a_j = 1$, $a_i = a$ and $a_r = b$ is a solution to the above equation. Choose the minimal positive a and b to get the primitive solution. □

Theorem 4.2. *If r is prime, then the orbits of finite subgroups of rank r are separated by*

$$\{v_i \mid i = 1, \dots, r-1\} \cup \{v_{ij} \mid 1 \le i < j < r\}.$$

Proof. Consider an arbitrary non-negative non-zero solution $(a_1, \dots, a_r)$ to Equation 4.1 and define $f := d_{1,r}^{a_1} \cdots d_{r-1,r}^{a_{r-1}} d_{r,r}^{-a_r}$. Suppose E is a rank r subgroup. If $f(E) = 0$, then $d_{i,r}(E) = 0$ for some $i < r$ and $f(E)$ is determined by $v_i(E)$. Thus we may assume $f(E) \neq 0$.

Since r is prime, it follows from Lemma 4.1, that the invariants v_{ij} exist. Thus we can write

$$f = v_{12}^{a_2} v_{13}^{a_3} \cdots v_{1(r-1)}^{a_{r-1}} d_{1,r}^a d_{r,r}^{-b}$$

for some, not necessarily positive, integers a and b. Since $f/(v_{12}^{a_2} v_{13}^{a_3} \cdots v_{1(r-1)}^{a_{r-1}})$ is invariant, $d_{1,r}^a d_{r,r}^{-b} = v_1^c$ for some integer c. If $d_{1,r}(E) \neq 0$, then $v_1^c(E)$ is

well-defined even if c is negative and $f(E)$ is determined by $v_1(E)$ and $v_{1j}(E)$ for $j > 1$.

Suppose $d_{1,r}(E) = 0$. Then, since $f(E) \neq 0$, we have $a_1 = 0$. Hence

$$f = v_{23}^{a_3} v_{24}^{a_4} \cdots v_{2(r-1)}^{a_{r-1}} v_2^a$$

for some, not necessarily positive, integer a. If $d_{2,r}(E) \neq 0$, then $v_2^a(E)$ is well-defined and $f(E)$ is determined by $v_2(E)$ and $v_{2j}(E)$ for $j > 2$.

Continuing in this fashion, the problem is reduced to $d_{i,r}(E) = 0$ for $i < r-1$ and $f(E) \neq 0$. Thus f has support $\{d_{r,r-1}, d_{r,r}^{-1}\}$ and is a positive power of v_{r-1}. □

Theorem 4.3. *The orbits of finite subgroups of rank* 4 *are separated by* $\{v_1, v_2, v_3, v_{12}.v_{13}, v_{32}\}$.

Proof. The proof is similar to the proof of Theorem 4.2. Using Lemma 4.1, the invariants v_{12}, v_{13} and v_{32} exist. The set $\{v_1, v_{12}, v_{13}\}$ separates orbits with $d_{1,4}(E) \neq 0$, the set $\{v_3, v_{32}\}$ separates orbits with $d_{1,4}(E) = 0$, $d_{3,4}(E) \neq 0$, and v_2 separates orbits with $d_{1,4}(E) = 0$, $d_{3,4}(E) = 0$ and $d_{2,4}(E) \neq 0$. □

Lemma 4.4. *Suppose the greatest common divisor of* r *and* i *is* 2 *and* j *is odd. If* $d_{j,r}^{a_j} d_{i,r}^{a_i} d_{r,r}^{-a_r}$ *is a primitive invariant, then* $p+1$ *divides* a_j*. Furthermore, there exists a primitive invariant*

$$u_{ij} := d_{j,r}^{p+1} d_{i,r}^{a} d_{r,r}^{-b}.$$

Proof. To find invariants with support $\{d_{i,r}, d_{j,r}, d_{r,r}^{-1}\}$, we consider the equation

$$a_j p^{r-j}(p^j - 1) = a_r(p^r - 1) - a_i p^{r-i}(p^i - 1).$$

Since r and i are even, $p^2 - 1$ divides both $p^r - 1$ and $p^i - 1$. Hence $p^2 - 1$ divides $a_j p^{r-j}(p^j - 1)$. However, since j is odd, $p + 1$ is relatively prime to $p^j - 1$. Thus $p + 1$ divides a_j.

Since 2 is the greatest common divisor of r and i, there exist positive integers $\bar{a}$ and $\bar{b}$ with $p^2-1 = \bar{b}(p^r-1)-\bar{a}p^{r-i}(p^i-1)$. Note that $c := p^{r-j}(p^j-1)/(p-1)$ is a positive integer. Define $a := c\bar{a}$ and $b := c\bar{b}$. Then $(p+1)p^{r-j}(p^j-1) = b(p^r-1) - ap^{r-i}(p^i-1)$. Hence $a_j = p+1$, $a_i = a$ and $a_r = b$ is a solution to the above equation. Choose the minimal positive a and b to get the primitive solution. □

Theorem 4.5. *The orbits of finite subgroups of rank* 6 *are separated by* $\{v_1, v_2, v_3, v_4, v_5, v_{12}, v_{13}, v_{14}, v_{15}, v_{52}, v_{53}, v_{54}, v_{24}, u_{23}, u_{43}\}$.

Proof. The proof is similar to the proof of Theorem 4.2. Using Lemmas 4.1 and 4.4, the listed invariants exist. The set $\{v_1, v_{12}, v_{13}, v_{14}, v_{15}\}$ separates orbits with $d_{1,6}(E) \neq 0$. The set $\{v_5, v_{52}, v_{53}, v_{54}\}$ separates orbits with $d_{1,6}(E) = 0$ and $d_{5,6}(E) \neq 0$. The set $\{v_2, v_{24}, u_{23}\}$ separates orbits with $d_{1,6}(E) = 0$, $d_{5,6}(E) = 0$ and $d_{2,6}(E) \neq 0$. The set $\{v_4, u_{43}\}$ separates orbits with $d_{1,6}(E) = 0$, $d_{5,6}(E) = 0$, $d_{2,6}(E) = 0$ and $d_{4,6}(E) \neq 0$. Finally, v_3 separates orbits with $d_{1,6}(E) = 0$, $d_{5,6}(E) = 0$, $d_{2,6}(E) = 0$, $d_{4,6}(E) = 0$ and $d_{3,6}(E) \neq 0$ □

Theorem 4.6. *The orbits of finite subgroups of rank* 8 *are separated by*

$$\{v_i \mid 1 \leq i < 8\} \cup \{v_{ij} \mid i \,\text{odd}\,, i < j\} \cup \{v_{24}, v_{26}, v_{64}\}.$$

Proof. The proof is similar to the proof of Theorem 4.2. Using Lemma 4.1, the listed invariants exist. We then "eliminate" variables in the following order: $1, 3, 5, 7, 2, 6, 4$. □

Theorem 4.7. *The orbits of finite subgroups of rank* 9 *are separated by*

$$\{v_i \mid 1 \leq i < 9\} \cup \{v_{ij} \mid i \notin \{3, 6\}\,, i < j\} \cup \{v_{36}\}.$$

Proof. The proof is similar to the proof of Theorem 4.2. Using Lemma 4.1, the listed invariants exist. We then "eliminate" variables in the following order: $1, 2, 4, 5, 7, 8, 3, 6$. □

Theorem 4.8. *The orbits of finite subgroups of rank* 10 *are separated by*

$$\{v_i \mid 1 \leq i < 10\} \cup \{v_{ij} \mid i \in \{1, 3, 7, 9\}\,, i < j\} \cup \{v_{ij}, u_{i5} \mid i, j \,\text{even},\ i < j\}.$$

Proof. The proof is similar to the proof of Theorem 4.2. Using Lemmas 4.1 and 4.4, the listed invariants exist. We then "eliminate" variables in the following order: $1, 3, 7, 9, 2, 4, 6, 8, 5$. □

5. Groups of Rank 3

In this section we consider finite subgroups of rank 3. It follows from Theorem 4.2 that the orbits of rank 3 subgroups are separated by v_1, v_2 and v_{12}. To construct a generating set for the invariants we add the intersection of the integer lattice with the line segment joining v_2 and v_{12}

Theorem 5.1. $\mathbb{F}[d_{1,3}, d_{2,3}, d_{3,3}^{-1}]^{\mathbb{F}^*}$ *is generated by* v_1, v_{12}, v_2 *and*

$$f_i = {d_{1,3}}^{p-i} {d_{2,3}}^{i(p+1)+1} {d_{3,3}}^{-(i+1)p}$$

for $i = 1, 2, \ldots, p-1$.

Proof. In this case Equation 4.1 becomes $a_1p^2 + a_2(p^2+p) = a_3(p^2+p+1)$. We will work with exponent sequences so $v_1 = (p^2+p+1, 0, p^2)$, $v_{12} = (p, 1, p)$, $v_2 = (0, p^2+p+1, p^2+p)$ and

$$f_i = v_{12} + \frac{i}{p}(v_2 - v_{12}) = (p, 1, p) + i(-1, p+1, p) = (p-i, (i+1)p+1, (i+1)p).$$

From this it is not hard to see that the primitive solutions of height one are precisely v_2, v_{12} and f_i for $i = 1, \ldots, p-1$. Note that v_1 has height $p+1$. We will show that there are no additional primitive solutions. Suppose $Q = (a_1, a_2, a_3)$ is an additional primitive solution. If $a_1 = 0$, then Q is a multiple of v_2, and if $a_2 = 0$, then Q is a multiple v_1. Thus we may assume $a_1 \geq 1$ and $a_2 \geq 1$. If $a_1 \geq p$, then $Q - v_{12}$ is a non-negative solution, so we may assume $p > a_1 \geq 1$. We may also assume that the height of Q is at least two.

Since p divides a_3, we can write $a_3 = p\widetilde{a}_3$ with $\widetilde{a}_3$ a positive integer. Hence the equation becomes

$$a_1 p + a_2(p+1) = \widetilde{a}_3(p^2+p+1).$$

Write $a_1 = p - i$ for $i \in \{1, \ldots, p-1\}$. We will show that $Q - f_i$ is non-negative, contradicting the assumption that Q was an additional primitive solution. Observe that $\widetilde{a}_3 \equiv i+1 \pmod{p+1}$. Since $i+1 \in \{2, \ldots, p\}$ and $\widetilde{a}_3$ is positive, this means $\widetilde{a}_3 \geq i+1$ and $a_3 \geq p(i+1)$. Since the height is at least two,

$$a_2 \geq 2 + a_3 - a_1 = a_3 + i - p + 2 \geq p(i+1) + i - p + 2 = (p+1)i + 2.$$

Hence $Q - f_i$ is non-negative, as required. □

If V is a subgroup of rank 3, then $\lambda(V) = (3)$, or $\lambda(V) = (2,1)$ or $\lambda(V) = (1,1,1)$. Applying Theorems 3.2 and 3.4 gives the following.

Theorem 5.2. *Let V be a rank 3 subgroup of $(\mathbb{F}, +)$. Then*

(i) $\lambda(V) = (2,1)$ *if and only if* $v_{12}(V) = 1$.
(ii) $\lambda(V) = (3)$ *if and only if* $d_{1,3}(V) = d_{2,3}(V) = 0$ *and* $d_{3,3}(V) \neq 0$.

6. Groups of Rank 4

In this section we consider finite subgroups of rank 4. It follows from Theorem 4.3 that the orbits of rank 4 subgroups are separated by v_1, v_2, v_3, v_{12}, v_{13} and v_{32}. To extend this to a generating set, we need to identify the primitive solutions to the equation

$$a_1(p^4 - p^3) + a_2(p^4 - p^2) + a_3(p^4 - p) = a_4(p^4 - 1). \tag{6.1}$$

Using Lemma 4.4, one additional solution is given by u_{23}. It will be convenient to work with the exponent sequences:

$$v_1 = \left(\frac{p^4-1}{p-1}, 0, 0, p^3\right),\ v_2 = \left(0, p^2+1, 0, p^2\right),\ v_3 = \left(0, 0, \frac{p^4-1}{p-1}, \frac{p(p^3-1)}{p-1}\right),$$
$$v_{12} = (p^2+p, 1, 0, p^2),\ v_{13} := (p, 0, 1, p),\ u_{23} = (0, p^2, p+1, p^2+p).$$

To generate the ring of invariants, we need the following additional solutions:

(i) lattice points on the line segment joining v_2 to v_{12}:

$$\mathcal{L} := \{((p+1)j, p^2 + 1 - jp, 0, p^2) \mid 0 \leq j \leq p\};$$

(ii) lattice points in the triangle with vertices v_{13}, v_3 and u_{23}:

$$\Delta := \{iv_{13}/p + ju_{23}/p^2 + (p^2 - pi - j)v_3/p^2 \mid 0 \leq i \leq p, 0 \leq j \leq p^2 - ip\}.$$

Note that

$$\begin{aligned} iv_{13}/p &+ ju_{23}/p^2 + (p^2 - pi - j)v_3/p^2 \\ &= (i, j, p^3 + (1-i)p^2 + (1-i-j)(p+1), p^3 + (1-i)p^2 + (1-i-j)p) \end{aligned}$$

and v_{32} is given by taking $i = 0$ and $j = 1$.

Theorem 6.1. *$\mathbb{F}[\mathbb{F}^5]^{GL_4(\mathbb{F}_p)\times\mathbb{F}^*}$ is generated as an $\mathbb{F}$-algebra by $d_{4,4}t$ and*

$$\{d_{1,4}^{a_1}d_{2,4}^{a_2}d_{3,4}^{a_3}t^{a_4} \mid (a_1,a_2,a_3,a_4) \in \{v_1\}\cup\mathcal{L}\cup\Delta\}.$$

Proof. We prove the theorem by identifying the primitive solutions to Equation 6.1. Suppose, by way of contradiction $Q = (a_1, a_2, a_3, a_4)$ is a primitive solution not included in the above list.

We start by considering solutions with $a_3 = 0$. In this case p^2 divides a_4 and $p + 1$ divides a_1. Writing $a_4 = \bar{a}_4p^2$ and $a_1 = \bar{a}_1(p + 1)$, Equation 6.1 becomes $\bar{a}_1p + a_2 = \bar{a}_4p^2$. If $a_2 = 0$, then Q is a multiple of v_1. If $a_2 > 0$ and $\bar{a}_1 \geq p$, then $Q - v_{12}$ is non-negative. Thus we may assume $\bar{a}_1 < p$. It is then easy to see that $Q - u$ is non-negative for some $u \in \mathcal{L}$.

In the following, we assume $a_3 \neq 0$. It is clear that p divides a_4; writing $a_4 = p\widetilde{a}_4$, Equation 6.1 becomes

$$a_1p^2 + a_2(p^2 + p) + a_3(p^2 + p + 1) = \widetilde{a}_4(p^3 + p^2 + p + 1).$$

Thus p divides $a_3 - \widetilde{a}_4$; writing $a_3 - \widetilde{a}_4 = p\widetilde{a}_3$, gives

$$a_1p + a_2(p + 1) + \widetilde{a}_3(p^2 + p + 1) = \widetilde{a}_4p^2.$$

Hence p divides $a_2 + \widetilde{a}_3$; writing $a_2 + \widetilde{a}_3 = p\widetilde{a}_2$, gives

$$a_1 + \widetilde{a}_2(p + 1) + \widetilde{a}_3p = \widetilde{a}_4p.$$

Finally, writing $a_1 + \widetilde{a}_2 = p\widetilde{a}_1$, gives

$$\widetilde{a}_1 + \widetilde{a}_2 + \widetilde{a}_3 = \widetilde{a}_4.$$

Note that $\widetilde{a}_1$ is in fact the height of the solution. In this new basis, the constraints $a_1 \geq 0$, $a_2 \geq 0$, $a_3 > 0$ and $a_4 > 0$ become

$$p\widetilde{a}_1 \geq \widetilde{a}_2, \qquad p\widetilde{a}_2 \geq \widetilde{a}_3, \qquad p\widetilde{a}_3 > -\widetilde{a}_4, \qquad \widetilde{a}_4 > 0.$$

Substituting and simplifying gives $a_1 + a_3 = (p + 1)(\widetilde{a}_1 + \widetilde{a}_3)$. Since we are assuming $a_3 > 0$, we have $\widetilde{a}_1 + \widetilde{a}_3 > 0$. Suppose Q has height one, in other words $\widetilde{a}_1 = 1$. Then $\widetilde{a}_3 \geq 0$. Since $p\widetilde{a}_1 \geq \widetilde{a}_2$ and $p\widetilde{a}_2 \geq \widetilde{a}_3$, we conclude $\widetilde{a}_2 \in \{0, \ldots, p\}$ and $\widetilde{a}_3 \in \{0, \ldots, p\widetilde{a}_2\}$. The (i, j) solution from the family Δ gives a solution with $\widetilde{a}_2 = p - i$ and $\widetilde{a}_3 = p^2 - pi - j$. Thus Δ is precisely the set of primitive solutions with height one and $a_3 > 0$.

Suppose $\widetilde{a}_1 > 1$ and $a_3 > 0$. If $a_1 \geq p$, then $Q - v_{13}$ is non-negative, so we may assume $a_1 < p$. Thus $p\widetilde{a}_1 - \widetilde{a}_2 \leq p - 1$, giving $\widetilde{a}_2 \geq p\widetilde{a}_1 - p + 1 \geq 2p - p + 1 = p + 1$. Hence $a_2 \geq p^2 + p - \widetilde{a}_3$. If $a_2 > p^2 + 1$ then $Q - v_2$ is non-negative, so we may assume $\widetilde{a}_3 \geq p$. Let u denote the solution in Δ given by $i = a_1$ and $j = p^2 - pa_1$. Then $u = (a_1, p^2 - pa_1, p + 1 - a_1, p^2 + p - pa_1)$. Since $a_3 = (p + 1)(\widetilde{a}_1 + \widetilde{a}_3) - a_1 > (p + 1) - a_1$, we have $Q - u$ is non-negative unless $a_2 < p^2 - pa_1$. Thus we may assume $p^2 - 1 \geq pa_1 + a_2$. Thus $p^2-1 \geq p^2\widetilde{a}_1-\widetilde{a}_3 \geq 2p^2-\widetilde{a}_3$, giving $\widetilde{a}_3 \geq p_1^2$. Hence $\widetilde{a}_4 = \widetilde{a}_1+\widetilde{a}_2+\widetilde{a}_3 \geq pr+p+3$, giving $a_3 = \widetilde{a}_4 + p\widetilde{a}_3 \geq p^3 + p^2 + 2p + 3$. Therefore $Q - v_3$ is non-negative, giving the required contradiction. □

Theorem 6.2. *Suppose E is a subgroup of rank 4. Then E contains a dilation of $\mathbb{F}_{p^2}$ if and only if*

$$\left(d_{1,4}^{p^2}d_{2,4}^{p}d_{3,4} - d_{4,4}^{p}d_{1,4}^{p^2+1} - d_{3,4}^{p^2+1}\right)(E) = 0.$$

Proof. The argument is similar to the proof of Theorem 3.4. The subgroup $E = \mathrm{Span}_{\mathbb{F}_P}\{c_1, c_2, c_3, c_4\}$ contains a dilation of $\mathbb{F}_{p^2}$ if and only if there is a $g \in GL_4(\mathbb{F}_p)$ such that $d_{1,2}g(c_1, c_2, c_3, c_4) = 0$. The stabiliser of $d_{1,2}$ has order $|GL_2(\mathbb{F}_p)|^2p^4$. Therefore the $GL_4(\mathbb{F}_p)$-orbit product of $d_{1,2}$ has order

$$\deg(d_{1,2})(p^2+1)(p^3-1)/(p-1) = p(p^2+1)(p^3-1).$$

The polynomial

$$H := d_{1,4}^{p^2}d_{2,4}^{p}d_{3,4} - d_{4,4}^{p}d_{1,4}^{p^2+1} - d_{3,4}^{p^2+1}$$

is a non-zero homogeneous $GL_4(\mathbb{F}_p)$-invariant of degree $p(p^2+1)(p^3-1)$. We will show that $d_{1,2}$ divides H and, therefore, H is, up to multiplication by a non-zero scalar, the $GL_4(\mathbb{F}_p)$-orbit product of $d_{1,2}$. Hence $H(E) = 0$ if and only if E contains a dilation of $\mathbb{F}_{p^2}$.

It follows from Theorem 3.4 that $d_{1,2}$ divides $d_{1,3}^{p}d_{2,3} - d_{3,3}^{p}$. Thus it is sufficient to work modulo the ideal $\mathfrak{r} := \langle d_{1,3}^{p}d_{2,3} - d_{3,3}^{p}\rangle$. We set $h = D_3^{p-1}(x_4)$ and use the equations $d_{i,4} = d_{i,3}^{p} - d_{i-1,3}h$. Substituting and simplifying gives $d_{2,4}^{p}d_{3,4} - d_{4,4}^{p}d_{1,4} \equiv_{\mathfrak{r}} d_{2,3}^{p^2}d_{3,4}$ and $d_{1,4}d_{2,3} - d_{3,4} \equiv_{\mathfrak{r}} 0$. Thus

$$H = d_{1,4}^{p^2}\left(d_{2,4}^{p}d_{3,4} - d_{4,4}^{p}d_{1,4}\right) - d_{3,4}^{p^2+1} \equiv_{\mathfrak{r}} d_{3,4}\left(d_{1,4}d_{2,3} - d_{3,4}\right)^{p^2} \equiv_{\mathfrak{r}} 0.$$

□

Theorem 6.3. *Suppose E is a subgroup of rank 4. Then*

(i) $\lambda(E) = (4)$ *if and only if* $d_{i,4}(E) = 0$ *for* $i \in \{1, 2, 3\}$;
(ii) $\lambda(E) = (2, 2)$ *if and only if* $d_{1,4}(E) = d_{3,4}(E) = 0$ *and* $d_{2,4}(E) \neq 0$;
(iii) $\lambda(E) = (3, 1)$ *if and only if* $d_{2,4}(E) = 0$ *and* $v_{13}(E) = 1$;
(iv) $\lambda(E) = (2, 1, 1)$ *if and only if* $d_{1,4}(E) \neq 0$, $d_{2,4}(E) \neq 0$ *and* $\left((v_{12} - v_{13})^p v_{13} - v_1\right)(E) = 0$.

Proof. Parts (i) and (ii) follow from Theorem 3.2. Part (iii) follows from Theorem 3.4. For part (iv), if $\lambda(E) = (2, 1, 1)$ then E contains a dilation of $\mathbb{F}_{p^2}$ but does not contain a dilation of $\mathbb{F}_{p^3}$ and is not an $\mathbb{F}_{p^2}$-subspace. Therefore, using Theorem 6.2, $\lambda(E) = (2, 1, 1)$ if and only if $H(E) = 0$ but $d_{2,4}(E) \neq 0$ and $d_{1,4}(E) \neq 0$. Multiplying H by $d_{1,4}^{p^3+p}/d_{4,4}^{p^3+1}$ gives $(v_{12} - v_{13})^p v_{13} - v_1$. □

Remark 6.4. *It is not clear that the partition is the best way to understand the structure of a subgroup. Consider $V = \mathbb{F}_{p^3} + \mathbb{F}_{p^2} \subset \overline{\mathbb{F}_p}$. This is a rank 4 subgroup with partition $(3, 1)$. The equivalence class of the subgroup is determined by $v_{12}(V) = 0$, $v_{13}(V) = 1$ and $v_1(V) = -1$.*

7. Invariants for Subgroups of Rank 5

In this section we will construct a generating set for $\mathbb{F}[\mathbb{F}^6]^{GL_5(\mathbb{F}_p)\times\mathbb{F}^*}$. To do this, we identify the primitive non-negative integer solutions to

$$a_1p^4+a_2p^3\left(\frac{p^2-1}{p-1}\right)+a_3p^2\left(\frac{p^3-1}{p-1}\right)+a_4p\left(\frac{p^4-1}{p-1}\right)=a_5\left(\frac{p^5-1}{p-1}\right). \tag{7.1}$$

Since we are interested in non-zero solutions, we assume $a_5>0$. As in the proof of Theorem 6.1, it will be useful to introduce an alternative basis: $a_5=p\widetilde{a}_5$, $a_4=p\widetilde{a}_4+\widetilde{a}_5$, $a_3=p\widetilde{a}_3-\widetilde{a}_4$, $a_2=p\widetilde{a}_2-\widetilde{a}_3$ and $a_1=p\widetilde{a}_1-\widetilde{a}_2$. In these new variables, the above equation becomes $\widetilde{a}_1+\widetilde{a}_2+\widetilde{a}_3+\widetilde{a}_4=\widetilde{a}_5$, the height of the solution, $a_1+a_2+a_3+a_4-a_5$, becomes $\widetilde{a}_1$ and the constraints $a_5>0$ and $a_i\geq 0$ for $i\in\{1,2,3,4\}$ become

$$p\widetilde{a}_1\geq\widetilde{a}_2,\quad p\widetilde{a}_2\geq\widetilde{a}_3,\quad p\widetilde{a}_3\geq\widetilde{a}_4,\quad p\widetilde{a}_4\geq-\widetilde{a}_5,\quad \widetilde{a}_5>0.$$

We will write $(a_1,a_2,a_3,a_4,a_5)=[\widetilde{a}_1,\widetilde{a}_2,\widetilde{a}_3,\widetilde{a}_4,\widetilde{a}_5]$. It follows from Theorem 4.2 that orbits of finite subgroups of rank 5 are separated by

$$\{v_1,v_2,v_3,v_4,v_{12},v_{13},v_{14},v_{23},v_{24},v_{34}\}.$$

We will extend this set to a generating set. Define

$$w_{24}:=(0,p^2,0,p+1,p^2+p)=[1,p,0,0,p+1]$$

and

$$w_{34}:=(0,0,p^3,p^2+p+1)=[1,p,p^2,0,p^2+p+1].$$

Let $\mathcal{T}_1$ denote the lattice points in the tetrahedron with vertices v_{14}, v_4, w_{24}, w_{34}. Define

$$w_{43}:=(0,0,p^3+1,p^2,p^3+p^2)=[1,p,p^2,-1,p^2+p].$$

Let $\mathcal{T}_2$ denote the lattice points in the tetrahedron with vertices v_{14}, v_{23}, v_{34}, w_{43}. Define

$$v_{312}=(p-1,1,p^2-p+1,0,p^2)=[1,1,p-1,-1,p].$$

Let Δ_1 denote the lattice points in the triangle with vertices v_{32}, v_{23}, v_{312}. Define

$$v_{123}:=(p^2-1,p,1,0,p^2)=[p,1,0,-1,p].$$

Let Δ_2 denote the lattice points in the triangle with vertices v_{312}, v_{23}, v_{123}. Define

$$w_{21}:=(p,p^3+1,0,0,p^3)=[p+1,p^2,-1,-p,p^2]$$

and

$$w_{24}:=(0,p^3+p^2+1,0,p,p^3+p^2)=[p+1,p^2+p,-1,-p,p^2+p].$$

Let Δ_3 denote the lattice points in the triangle with vertices v_2, w_{21}, w_{24}. Let $\mathcal{L}_1$ denote the lattice points on the line segment joining v_3 to v_{13}. Let $\mathcal{L}_2$ denote the lattice points on the line segment joining v_{12} to w_{21}. Define

$$\mathcal{S}:=\left(\bigcup_{i=1}^{2}\mathcal{T}_i\right)\cup\left(\bigcup_{i=1}^{3}\Delta_i\right)\cup\left(\bigcup_{i=1}^{2}\mathcal{L}_i\right)\cup\{v_1\}.$$

Theorem 7.1. $\mathbb{F}[\mathbb{F}^6]^{GL_5(\mathbb{F}_p)\times\mathbb{F}^*}$ *is generated as an* $\mathbb{F}$*-algebra by* $d_{5,5}t$ *and*

$$\{d_{1,4}^{a_1}d_{2,4}^{a_2}d_{3,4}^{a_3}d_{4,4}^{a_4}t^{a_5} \mid (a_1,a_2,a_3,a_4,a_5)\in\mathcal{S}\}.$$

The remainder of this section is dedicated to the proof of Theorem 7.1

Lemma 7.2. *The elements in* $\mathcal{T}_1$ *are precisely the solutions with height one and* $\widetilde{a}_4 \geq 0$*.*

Proof. Note that if $\widetilde{a}_4 \geq 0$ then $a_4 > 0$. For $i \in \{0,\dots,p\}$, $j \in \{0,\dots,p^2-ip\}$ and $k \in \{0,\dots,p^3-ip^2-jp\}$, define

$$u_{ijk} := \frac{i}{p}v_{14} + \frac{j}{p^2}w_{24} + \frac{k}{p^3}w_{34} + \left(1-\frac{i}{p}-\frac{j}{p^2}-\frac{k}{p^3}\right)v_4.$$

Define $c_{ijk} := p^4+(1-i)p^3+(1-i-j)p^2+(1-i-j-k)p$. Simplifying gives

$$u_{ijk} = (i,j,k,c_{ijk}+(1-i-j-k),c_{ijk})\,.$$

Write $\widetilde{i} := p-i$, $\widetilde{j} := p^2-ip-j$, $\widetilde{k} := p^3-p^2i-pj-k$. Then

$$u_{ijk} = [1,\widetilde{i},\widetilde{j},\widetilde{k},1+\widetilde{i}+\widetilde{j}+\widetilde{k}]$$

with $p \geq \widetilde{i}$, $p\widetilde{i} \geq j$ and $p\widetilde{j} \geq \widetilde{k} \geq 0$. From this it is clear that $\mathcal{T}_1$ consists of the solutions with height one and $\widetilde{a}_4 \geq 0$. □

Lemma 7.3. *All solutions with height one,* $a_4 > 0$ *and* $\widetilde{a}_4 < 0$ *lie in* $\mathcal{T}_2$*.*

Proof. Substituting and simplifying gives $a_1+a_4 = (p+1)(\widetilde{a}_1+\widetilde{a}_4)+\widetilde{a}_3$. Since we are assuming $a_4 > 0$, $\widetilde{a}_4 < 0$, and $\widetilde{a}_1 = 1$, we conclude that $\widetilde{a}_3 > 0$. Using the basic constraints and the assumption $\widetilde{a}_1 = 1$, gives $p \geq \widetilde{a}_2 > 0$ and $p^2 \geq p\widetilde{a}_2 \geq \widetilde{a}_3$. Since $p\widetilde{a}_4 = a_4-\widetilde{a}_5 \geq 1-\widetilde{a}_5 = 1-(\widetilde{a}_1+\widetilde{a}_2+\widetilde{a}_3+\widetilde{a}_4) = -\widetilde{a}_2-\widetilde{a}_3-\widetilde{a}_4$, we have $\widetilde{a}_2+\widetilde{a}_3 \geq -(p+1)\widetilde{a}_4$. Thus $p(p+1) \geq -\widetilde{a}_4(p+1)$ and $-p \leq \widetilde{a}_4 \leq -1$. Define

$$\begin{aligned} t_{ijk} := \left(\frac{i}{p}-\frac{j}{p^2}\right)v_{23} &- \left(\frac{k}{p-1}+\frac{i}{p(p-1)}\right)v_{34} \\ &+ \left(\frac{k}{p-1}+\frac{i}{p(p-1)}+\frac{j}{p^2}\right)w_{43} + \left(1-\frac{i}{p}\right)v_{14} \end{aligned}$$

for $p \geq i \geq 0$, $pi \geq j \geq 0$ and $-1 \geq k \geq -p$. Note that, as long as $i+j+(p+1)k \geq 0$, the coefficients are non-negative and $t_{ijk} \in \mathcal{T}_2$. Substituting and simplifying gives

$$\begin{aligned} t_{ijk} &= (p-i,pi-j,pj-k,1+i+j+k(p+1),p(1+i+j+k)) \\ &= [1,i,j,k,1+i+j+k]. \end{aligned}$$

Thus every solution with height one, $a_4 > 0$, $\widetilde{a}_4$ and $\widetilde{a}_1 > 1$ lies in $\mathcal{T}_2$. □

Lemma 7.4. *The solutions with height one and* $a_4 = 0$ *lie in* Δ_1*.*

Proof. Arguing as in the proof of Lemma 7.3, we have $p^2 \geq p\widetilde{a}_2 \geq \widetilde{a}_3 > 0$. Since $-p\widetilde{a}_4 = \widetilde{a}_5 - a_4 = \widetilde{a}_5 = 1 + \widetilde{a}_2 + \widetilde{a}_3 + \widetilde{a}_4$, we have $\widetilde{a}_2 + \widetilde{a}_3 + 1 = -(p+1)\widetilde{a}_4$. Writing $p\widetilde{a}_2 - \widetilde{a}_3 = k$, this equation becomes $(p+1)\widetilde{a}_2 + 1 - k = -(p+1)\widetilde{a}_4$. Thus $k \equiv 1 \pmod{p+1}$. Define

$$t_{ij} := \frac{i}{p-1}v_{23} + \frac{j}{p-1}v_{312} + \frac{p-1-i-j}{p-1}v_{32}$$

for $i, j \in \{0, \ldots, p-1\}$ with $i + j \leq p-1$. Substituting and simplifying gives

$$\begin{aligned} t_{ij} &= \big(j, i(p+1)+1, p^3 - j(p^2+1) - i(p^2+p+1), 0, p^2(p-i-j)\big) \\ &= [1, p-j, p^2 - 1 - jp - i(p+1), -(p-i-j), p(p-i-j)]. \end{aligned}$$

Thus t_{ij} lies in Δ_1. We may choose $j \in \{0, \ldots, p-1\}$ so that $\widetilde{a}_2 = p - j$ and choose $i \in \{0, \ldots, p-1-j\}$ so that $i(p+1)+1 = k$ with $k = p\widetilde{a}_2 - \widetilde{a}_3$. Our solution is given by t_{ij} for these choices. □

Lemma 7.5. *There are no primitive solutions with $\widetilde{a}_1 > 1$ and $\widetilde{a}_4 \geq 0$.*

Proof. Suppose, by way of contradiction, that $Q = (a_1, a_2, a_3, a_4, a_5)$ is a primitive solution with $\widetilde{a}_1 > 1$ and $\widetilde{a}_4 \geq 0$. Suppose $\widetilde{a}_4 \geq p^3$. Then, using the basic constraints, $\widetilde{a}_3 \geq p^2$ and $\widetilde{a}_2 > p$. Thus $Q - v_4$ is non-negative. Thus we may assume $\widetilde{a}_3 < p^3$. Similarly, using w_{34}, we may assume $\widetilde{a}_3 < p^2$ and using w_{24} we may assume $\widetilde{a}_2 < p$. Consider the primitive solutions u_{ijk} defined in the proof of Lemma 7.2. Taking $i = p - \widetilde{a}_2$, $j = p^2 - ip - \widetilde{a}_3$ and $k = p^3 - p^2 i - pj - \widetilde{a}_4$, gives a primitive solution with $Q - u_{ijk}$ non-negative. □

Remark 7.6. *Since $v_{23} = (0, p^2, 1, 0, p^2) = [1, p, 0, -1, p]$, the solutions in Δ_2 satisfy $\widetilde{a}_5 = p$, $\widetilde{a}_4 = -1$ and $a_4 = 0$. Define*

$$s_{ij} := \frac{i-1}{p-1}v_{23} + \frac{j-1}{p-1}v_{123} + \frac{p+1-(i+j)}{p-1}v_{312}$$

for $i, j \in \{1, \ldots, p\}$ and $i + j \leq p+1$. Then

$$\begin{aligned} s_{ij} &= \big(pi - j, (j-1)(p+1) + i, p\,(p+1-(i+j)) + 1, 0, p^2\big) \\ &= [i, j, p+1-(i+j), -1, p] \end{aligned}$$

lies in Δ_2.

Remark 7.7. *Since $v_2 = (0, p^4 + p^3 + p^2 + p + 1, 0, 0, p^4 + p^3)$, we see that the solutions in Δ_3 satisfy $a_3 = 0$, $a_4 < p$ and $\widetilde{a}_4 \leq -p$. Define*

$$u_{ij} := \frac{i}{p}w_{21} + \frac{j}{p}w_{24} + \frac{p-i-j}{p}v_2$$

for $i, j \in \{0, \ldots, p\}$ with $i + j \leq p$. Substituting and simplifying gives

$$\begin{aligned} u_{ij} &= (i, p^4 + p^2 + (p^3+1)(1-(i+j)) + p(1-i), 0, j, p^4 + p^3(1-(i+j)) + pj) \\ &= [p^2 + p + 1, p^3 + p^2 + p, -(p+1), -(p^2+p), p^3 + p^2] \\ &+ (i+j)[-p, -p^2, 1, p, -p^2] + [0, -i, 0, 0, j]. \end{aligned}$$

Hence $u_{ij} \in \Delta_3$.

Remark 7.8. *Since* $v_3 = (0, 0, p^4 + p^3 + p^2 + p + 1, 0, p^4 + p^3 + p^2)$ *and* $v_{13} = (p^2 + p, 0, 1, 0, p^2)$*, we see that the solutions in* $\mathcal{L}_1$ *satisfy* $a_2 = 0$*,* $a_4 = 0$ *and* $\widetilde{a}_4 < 0$*. Define* $u_i := (iv_{13} + (p^2 + p - i)v_3)/(p^2+p)$ *for* $i \in \{0, \ldots, p^2+p\}$*. Simplifying gives* $u_i = (i, 0, (p^2 + p - i)(p^2 + p) + 1, 0, (p^2 + p + 1 - i)p^2)$*.*

Remark 7.9. *Since* $v_{12} = (p^3 + p^2 + p, 1, 0, 0, p^3)$ *the solutions in* $\mathcal{L}_2$ *satisfy* $a_3 = 0$*,* $a_4 = 0$*,* $\widetilde{a}_4 = -p$ *and* $\widetilde{a}_5 = p^2$*. Examining* $v_{12} - w_{21}$*, we see that these solutions are given by*

$$(p^3 + p^2 + p - j(p + 1), 1 + jp, 0, 0, p^3) = [p^2 + p + 1 - j, j, -1, -p, p^2]$$

for $j \in \{0, \ldots, p^2\}$*.*

Suppose, by way of contradiction, that $Q = (a_1, a_2, a_3, a_4, a_5)$ is a primitive solution with $Q \notin \mathcal{S}$. Using Lemmas 7.2, 7.3, 7.4 and 7.5, we may assume $\widetilde{a}_1 > 1$ and $\widetilde{a}_4 < 0$.

We are left with four cases:

(i) $a_3 = 0$ and $a_4 = 0$;
(ii) $a_3 = 0$ and $a_4 > 0$;
(iii) $a_3 > 0$ and $a_4 = 0$;
(iv) $a_3 > 0$ and $a_4 > 0$.

Case (i). Suppose $a_3 = 0$, $a_4 = 0$, $\widetilde{a}_4 < 0$ and $\widetilde{a}_1 > 1$. If $a_2 = 0$, then Q is a multiple of v_1, so we may also assume $a_2 > 0$. Similarly, using v_2, we can assume $a_1 > 0$. Since $a_3 = a_4 = 0$, we have $a_5 = p\widetilde{a}_5 = -p^2\widetilde{a}_4 = -p^3\widetilde{a}_3$. Thus $\widetilde{a}_3 < 0$ and Equation 7.1 becomes $a_1p + a_2(p + 1) = -\widetilde{a}_3(p^4 + p^3 + p^2 + p + 1)$. Using $w_{21} = (p, p^3 + 1, 0, 0, p^3)$, we can assume that either $a_1 < p$ or $a_2 \leq p^3$.

Suppose $a_1 < p$. Write $a_1 = p - j$ (so $j \in \{1, \ldots, p - 1\}$). Then $a_2(p+1) = (j - p)p - \widetilde{a}_3(p^4 + p^3 + p^2 + p + 1)$. Note that $j + 1 \equiv -\widetilde{a}_3 \pmod{p+1}$. Write $-\widetilde{a}_3 = j + 1 + k(p + 1)$ (so $k \geq 0$). Thus

$$a_2(p+1) \geq (j-p)p+(j+1)(p^4+p^3+p^2+p+1) = (p+1)((j+1)(p^3+p)+j-p+1).$$

Hence $a_2 \geq (j + 1)(p^3 + p) + j - p + 1 = j(p^3 + p + 1) + p^3 + 1$. Within Δ_3, there are solutions on the line segment joining v_2 to w_{21} of the form $u_j = (p - j, p^3 + 1 + j(p^3 + p + 1), 0, 0, p^3 + jp^3))$ for $j \in \{0, \ldots, p\}$. Therefore $Q - u_j$ is non-negative and we may assume $a_2 \leq p^3$.

Suppose $p^3 \geq a_2$. Define $k \in \{1, \ldots, p\}$ so that $k \equiv a_2 \equiv -\widetilde{a}_3 \pmod{p}$. Note that $a_2 - k = jp$ for $j \in \{0, \ldots, p^2 - 1\}$ Then $a_1p \geq -(p+1)a_2 + k(p^4 + p^3 + p^2 + p + 1) = -pa_2 + k(p^4 + p^3 + p^2 + p) - jp$. Hence

$$a_1 \geq k(p^3 + p^2 + p + 1) - a_2 - j = k(p^3 + p^2 + p) - j(p + 1).$$

The family $\mathcal{L}_2$ includes a solution $u_j = (p^3 + p^2 + p - j(p + 1), 1 + jp, 0, 0, p^3)$ for $j \in \{0, \ldots, p^2\}$. Therefore $Q - u_j$ is non-negative, giving the required contradiction.

Case (ii). Suppose $a_3 = 0$, $a_4 > 0$, $\widetilde{a}_4 < 0$ and $\widetilde{a}_1 > 1$. Since $a_3 = 0$, we have $\widetilde{a}_4 = p\widetilde{a}_3$. Thus $\widetilde{a}_3 < 0$ and p divides $\widetilde{a}_4$. As in the proof of Lemma 7.3, substituting and simplifying gives $a_1 + a_4 = (p+1)(\widetilde{a}_1 + \widetilde{a}_4) + \widetilde{a}_3$. Since $a_4 > 0$ and $\widetilde{a}_3 < 0$, we have $\widetilde{a}_1 + \widetilde{a}_4 > 0$. Using $v_{14} = (p, 0, 0, 1, p)$, we may assume $a_1 < p$. Substituting and simplifying give $a_2 = p^2\widetilde{a}_1 - pa_1 - \widetilde{a}_3$. Since $\widetilde{a}_1 > -\widetilde{a}_4 = -p\widetilde{a}_3$, we have $a_2 \geq p^2(1-p\widetilde{a}_3) - \widetilde{a}_3 - pa_1 = (-\widetilde{a}_3)(p^3+1) + p^2 - pa_1$.

Suppose $a_1 + a_4 \geq p$. Let v denote the solution $u_{ij} \in \Delta_3$, as defined in Remark 7.7, with $i = a_1$ and $j = p - i$. Then $v = (i, p^3 + p^2 + 1 - ip, 0, p - i, p^3 + p(p-i))$. Since $-\widetilde{a}_3 \geq 1$, we have $a_2 \geq p^3 + p^2 + 1 - ip$. Thus $Q - v$ is non-negative.

Suppose $a_1 + a_4 < p$. Note that $a_1 + a_4 - \widetilde{a}_3$ is a positive multiple of $p+1$. Thus $-\widetilde{a}_3 \geq p + 1 - (a_1 + a_4)$ and $a_2 \geq (p^3+1)(p+1-(a_1+a_4)) + p^2 - pa_1 = p^4 + p^3 + p^2 p + 1 - (p^3+1)(a_1+a_4) - pa_1$. Thus $Q - v$ is non-negative where v is the solution $u_{ij} \in \Delta_3$ with $i = a_1$ and $j = a_4$.

Case (iii). Suppose $a_3 > 0$, $a_4 = 0$, $\widetilde{a}_4 < 0$ and $\widetilde{a}_1 > 1$. Since $a_4 = 0$, we have $a_5 = -p^2\widetilde{a}_4$. Using v_{13}, we may assume $a_1 < p^2 + p$ and using v_{23} we may assume $a_2 < p^2$. Note that $(p+1)(\widetilde{a}_2 + \widetilde{a}_4) + \widetilde{a}_1 = a_2 + \widetilde{a}_1 + \widetilde{a}_2 + \widetilde{a}_3 + (p+1)\widetilde{a}_4 = a_2$. Thus $(p+1)(\widetilde{a}_2 + \widetilde{a}_4) < a_2 < p^2$ and $\widetilde{a}_4 + \widetilde{a}_2 < p - 1$.

Further suppose $a_2 = 0$. Equation 7.1 becomes $a_1 p^4 + a_3 p^2(p^2 + p + 1) = a_5(p^4 + p^3 + p^2 + p + 1)$. Thus $a_1 + a_3 \equiv a_5 \pmod{p+1}$. Write $a_1 + a_3 - a_5 = k(p+1)$ with $k > 0$. Then the height of the solution is $\widetilde{a}_1 = a_1 + a_2 + a_3 + a_4 - a_5 = k(p+1)$. Hence $\widetilde{a}_2 = p\widetilde{a}_1 - a_1 = kp(p+1) - a_1$. Using $a_2 = 0$, gives $\widetilde{a}_3 = p\widetilde{a}_2 = kp^2(p+1) - a_1 p$. We have two expression for a_3:

$$a_3 = k(p+1) - a_1 + a_5 = k(p+1) - a_1 - p^2\widetilde{a}_4$$

and

$$a_3 = p\widetilde{a}_3 - \widetilde{a}_4 = p^2\widetilde{a}_2 - \widetilde{a}_4 = p^3 k(p+1) - p^2 a_1 - \widetilde{a}_4.$$

Equating these expressions and simplifying gives $a_1 - \widetilde{a}_4 = k(p^2+p+1)$. Using this to eliminate $\widetilde{a}_4$ from the second expression for a_3 give $a_3 = p^3 k(p+1) - p^2 a_1 + k(p^2+p+1) - a_1$. Let v denote the solution $u_i \in \mathcal{L}_1$, as defined in Remark 7.8, with $i = a_1$. Since $k > 0$, we see that $Q - v$ is non-negative. Thus we may assume $a_2 > 0$. Furthermore, using v_{32}, we may assume $a_3 < p^3$.

Suppose $\widetilde{a}_3 < 0$. Then $p^2 > a_2 = p\widetilde{a}_2 - \widetilde{a}_3 > p\widetilde{a}_2$. Hence $\widetilde{a}_2 < p$. Since $1 \leq a_3 = p\widetilde{a}_3 - \widetilde{a}_4$, we have $-\widetilde{a}_4 \geq p + 1$. Thus $\widetilde{a}_1 = \widetilde{a}_5 - (\widetilde{a}_2 + \widetilde{a}_3 + \widetilde{a}_4) = (-\widetilde{a}_4)(p+1) - \widetilde{a}_2 - \widetilde{a}_3 \geq (p+1)^2 + 1 - \widetilde{a}_2 > p^2 + p + 2$. Hence $a_1 = p\widetilde{a}_1 - \widetilde{a}_2 > p^3 + p^2 + p$, contradicting our assumption that $a_1 < p^2 + p$. Thus we may assume $\widetilde{a}_3 \geq 0$. Furthermore, since $p\widetilde{a}_2 \geq \widetilde{a}_3$, we may assume $\widetilde{a}_2 \geq 0$.

Note that $\widetilde{a}_1 + \widetilde{a}_2 + \widetilde{a}_3 = \widetilde{a}_5 - \widetilde{a}_4 = (-\widetilde{a}_4)(p+1)$. Thus if $\widetilde{a}_4 = -1$, we have $\widetilde{a}_3 = p + 1 - (\widetilde{a}_2 + \widetilde{a}_3) \geq 0$ and $Q = s_{ij}$ for $i = \widetilde{a}_1$ and $j = \widetilde{a}_2$, with $s_{ij} \in \Delta_2$ as defined in Remark 7.6. Thus we may assume $\widetilde{a}_4 < 1$.

Suppose $a_1 = 0$. Then $\widetilde{a}_2 = p\widetilde{a}_1 \geq 2p$ and $a_2 \geq 2p^2 - \widetilde{a}_3$. Thus $\widetilde{a}_3 > p^2$. Substituting and simplifying gives $a_1 = (p+1)(\widetilde{a}_1 + \widetilde{a}_4) + \widetilde{a}_3$. Thus $\widetilde{a}_3 =$

$(p+1)(-\widetilde{a}_4-\widetilde{a}_1) \geq p(p+1)$. Hence $a_3 = p\widetilde{a}_3 - \widetilde{a}_4 \geq p^3+p^2+2$, contradicting the assumption that $a_3 < p$. Thus we may assume $a_1 > 0$.

Suppose $a_1 < p$. Let u denote the solution $s_{ij} \in \Delta_2$ with $i = 1$ and $j = p-a_1$. Then $u = (a_1, p^2 - a_1(p+1), pa_1+1, 0, p^2)$. Thus either $a_2 < p^2 - a_1(p+1)$ or $a_3 \leq pa_1$. First suppose $a_2 < p^2 - a_1(p+1)$. Then $pa_1 + a_2 \leq p^2 - a_1 - 1 \leq p^2 - 1$, which implies $p^2 - 1 \geq p^2\widetilde{a}_1 - \widetilde{a}_3 \geq 2p^2 - \widetilde{a}_3$. Thus $\widetilde{a}_3 \geq p^2+1$ and $a_3 > p^3 + p + 2$, contradicting the assumption $a_3 < p^3$. Suppose $a_3 \leq pa_1$. Then $p\widetilde{a}_3 - \widetilde{a}_4 \leq p^2 - p$, giving $p\widetilde{a}_3 \leq p^2 - p + \widetilde{a}_4 \leq p^2 - p - 2$. Thus $\widetilde{a}_3 \leq p-2$. Hence $a_2 = p\widetilde{a}_2 - \widetilde{a}_3 \geq p\widetilde{a}_2 - p + 2$. Since $a_2 < p^2$, this forces $\widetilde{a}_2 < p$. Thus $a_1 = p\widetilde{a}_1 - \widetilde{a}_2 > p\widetilde{a}_1 - p \geq p$, giving the required contradiction. Thus we may assume $a_1 \geq p$.

If $a_1 \geq p^2 - 1$ then using the solution $v_{123} = (p^2-1, p, 1, 0, p^2)$ we see that $a_2 < p$. Next using the solutions $(ip-1, i, p^2 - ip + 1, 0, p^2)$ from Δ_2 with $i = a_2$ we have $a_3 \leq p^2 - a_2 p$. This implies that $p^2 \geq pa_2 + a_3 = p^2\widetilde{a}_2 - \widetilde{a}_4$ and thus $\widetilde{a}_2 = 0$. However $\widetilde{a}_2 = 0$ implies $\widetilde{a}_3 = 0$. If $\widetilde{a}_2 = \widetilde{a}_3 = 0$, then $a_2 = 0$, giving a contradiction. Thus we must have $a_1 \leq p^2 - 2$ and so $p \leq a_1 \leq p^2 - 2$.

The solution $v_{312} = (p-1, 1, p^2 - p + 1, 0, p^2)$ shows that $a_3 \leq p^2 - p$, from which we derive that $\widetilde{a}_3 \leq p-2$. If $a_2 < p$ then $p - 1 \geq a_2 = p\widetilde{a}_2 - \widetilde{a}_3 \geq p\widetilde{a}_2 - p + 2$ which yields $\widetilde{a}_2 \leq 1$. Hence $\widetilde{a}_1 = -\widetilde{a}_4(p+1) - \widetilde{a}_2 - \widetilde{a}_3 \geq 2(p+1) - 1 - (p-2) = p+3$ and thus $a_1 \geq p(p+3) - 1 = p^2 + 3p - 1$. Since $a_1 \leq p^2 - 2$ we have a contradiction and thus $a_2 \geq p$.

Again consider the solutions $(ip-1, i, p^2 - ip + 1, 0, p^2)$ from Δ_2. Choose i so that $ip - 1 \leq a_1 \leq ip + p - 2$. Since $a_2 \geq p > i$, we have $a_3 \leq p^2 - ip \leq p^2 - a_1 + p - 2$ which implies $a_1 + a_3 \leq p^2 + p - 2$. Thus $p\widetilde{a}_1 - \widetilde{a}_2 + p\widetilde{a}_3 - \widetilde{a}_4 \leq p^2 + p - 2$ and $p(\widetilde{a}_1 + \widetilde{a}_3) \leq p^2 + p - 2 + \widetilde{a}_2 + \widetilde{a}_4 < p^2 + 2p - 3$ since $\widetilde{a}_2 + \widetilde{a}_4 < p - 1$. This yields $\widetilde{a}_1 + \widetilde{a}_3 \leq p+1$. Therefore $-\widetilde{a}_4(p+1) = \widetilde{a}_1 + \widetilde{a}_2 + \widetilde{a}_3 \leq (p+1) + (p+1-\widetilde{a}_4)$ which implies $-\widetilde{a}_4 p \leq 2p + 2$. Since $\widetilde{a}_4 \leq -2$ we must have $\widetilde{a}_4 = -2$. Thus $\widetilde{a}_1 + \widetilde{a}_2 + \widetilde{a}_3 = 2(p+1)$. Since $\widetilde{a}_1 + \widetilde{a}_3 \leq p+1$, this gives $\widetilde{a}_2 \geq p+1$. However $\widetilde{a}_2 < p - 1 - \widetilde{a}_4 = p+1$, giving the required contradiction.

Case (iv). Suppose $a_3 > 0$, $a_4 > 0$, $\widetilde{a}_4 < 0$ and $\widetilde{a}_1 > 1$. Using $v_{14} = (p, 0, 0, 1, p)$, we may assume $a_1 < p$. Using $v_{23} = (0, p^2, 1, 0, p^2)$, we may assume $a_2 < p^2$. Thus $\widetilde{a}_2 = p\widetilde{a}_1 - a_1 > p$. Write $\widetilde{a}_2 = \alpha(p+1) + \beta$ with $0 \leq \alpha < p$ and $0 \leq \beta \leq p$. Let u denote the solution $t_{ijk} \in \mathcal{T}_2$, as defined in the proof of Lemma 7.3, with $i = p$, $k = \alpha - p$ and $j = p^2 - \alpha(p+1)$. Then $u = (0, \alpha(p+1), p^3 + p - \alpha(p^2+p+1), 1, p^3 + p - \alpha p^2)$. Thus $Q - u$ is non-negative if $a_3 \geq p^3 + p - \alpha(p^2+p+1)$. Hence we may assume $a_3 < p^3 + p - \alpha(p^2+p+1) \leq p^3 + p - p\alpha(p+1) \leq p^3 + p - p(a_2 - \beta) \leq p^3 + p^2 + p - pa_2$. Therefore $p^3 + p^2 + p > a_3 + pa_2 = p^2\widetilde{a}_2 - \widetilde{a}_4 > p^2\widetilde{a}_2$. From this we get $p+1 \geq \widetilde{a}_2$. Since we also have $\widetilde{a}_2 > p$, we conclude that $\widetilde{a}_2 = p+1$.

Since $a_1 = p\widetilde{a}_1 - \widetilde{a}_2 = p\widetilde{a}_1 - p - 1$, $\widetilde{a}_1 > 1$ and $a_1 < p$, we have $a_1 = p-1$. Furthermore, $p^2 > a_2 = p\widetilde{a}_2 - \widetilde{a}_3 = p(p+1) - \widetilde{a}_3$ gives $\widetilde{a}_3 > p$. Thus $a_3 = p\widetilde{a}_3 - \widetilde{a}_4 > p^2 + p$. Let v denote the solution $t_{ijk} \in \mathcal{T}_2$ with $i = 1$, $j = p$

and $k=-1$. Then $v=(p-1,0,p^2+1,1,p^2+p)$ and $Q-v$ in non-negative, giving the required contradiction.

8. Partitioning Rank 5 Subgroups

Applying Theorems 3.2 and 3.4, gives the following.

Theorem 8.1. *Suppose E is a subgroup of rank 5. Then*

(i) $\lambda(E)=(5)$ *if and only if* $d_{i,5}(E)=0$ *for* $i\in\{1,2,3,4\}$;
(ii) $\lambda(E)=(4,1)$ *if and only if* $d_{i,5}(E)=0$ *for* $i\in\{2,3\}$ *and* $v_{14}(E)=1$.

We do not have a complete characterisations of the partitions of the rank 5 subgroups. However the following approach has lead to a range of conjectures which have been verified for small primes using Magma [1].

Use h to denote $D_4^{p-1}(x_5)$. Then $d_{i,5}=d_{i,4}^p-d_{i-1,4}h\in A:=\mathbb{F}[d_{1,4},\dots,d_{4,4},h]$. Thus $\mathbb{F}[d_{1,5},\dots,d_{5,5}]$ is a subalgebra of A. Using Theorem 3.4, to show that a rank 5 subgroup $E=\mathrm{Span}_{\mathbb{F}_p}\{c_1,\dots,c_5\}$ contains a dilation of $\mathbb{F}_{p^3}$, it is sufficient to find $g\in GL_5(\mathbb{F}_p)$ such that $d_{2,4}g(c_1,\dots,c_5)=0$ and

$$\left(d_{1,4}^p d_{3,4}-d_{4,4}^p\right)g(c_1,\dots,c_5)=0.$$

In other words, we want to consider the union of the varieties determined by $\langle d_{2,4}g,\left(d_{1,4}^p d_{3,4}-d_{4,4}^p\right)g\rangle$. This is equivalent to the variety determined by the intersection of these ideals. Computing this intersection is equivalent to computing

$$\langle d_{2,4},\left(d_{1,4}^p d_{3,4}-d_{4,4}^p\right)\rangle\cap\mathbb{F}[d_{1,5},\dots,d_{5,5}].$$

This can be done computationally as an elimination of variables. Using this approach, the following has been verified for primes less than or equal to 13.

Conjecture 8.2. *A rank 5 subgroup E contains a dilation of $\mathbb{F}_{p^3}$ if and only if E is in the variety determined by $d_{4,5}^p-d_{3,5}d_{1,5}^p$, $d_{5,5}^p d_{1,5}-d_{4,5}d_{2,5}^p$ and $d_{5,5}^{ip}d_{4,5}^{p-i}-d_{3,5}d_{2,5}^{ip}d_{1,5}^{p-i}$ for $i\in\{1,\dots,p\}$.*

If E is in the variety determined by $\langle d_{1,4},d_{3,4}\rangle\cap\mathbb{F}[d_{1,5},\dots,d_{5,5}]$, then E contains a rank 4 subgroup which is a vector space over $\mathbb{F}_{p^2}$.

Conjecture 8.3. $\langle d_{1,4},d_{3,4}\rangle\cap\mathbb{F}[d_{1,5},\dots,d_{5,5}]=\langle d_{5,5}^p-d_{4,5}d_{1,5}^p,d_{3,5}^p-d_{2,5}d_{1,5}^p\rangle$.

Conjecture 8.3 has been verified for primes less than or equal to 17.

If E has partition $(4,1)$ then E is in the variety determined by

$$\langle d_{5,5}^p-d_{4,5}d_{1,5}^p,d_{3,5},d_{2,5}\rangle\subset\langle d_{5,5}^p-d_{4,5}d_{1,5}^p,d_{3,5}^p-d_{2,5}d_{1,5}^p\rangle.$$

Thus we expect a subgroup with partition $(2,2,1)$ to satisfy $d_{2,5}(E)\neq 0$.

Since $d_{5,5}(E)$ is non-zero for any rank 5 subgroup, we can replace $d_{5,5}^p-d_{4,5}d_{1,5}^p$ with $v_{14}-1$. If $(v_{14}-1)(E)=0$, then $d_{1,5}(E)\neq 0$ and so we can replace $d_{3,5}^p-d_{2,5}d_{1,5}^p$ with $v_{13}^p-v_{12}$. Thus

$$\langle d_{5,5}^p-d_{4,5}d_{1,5}^p,d_{3,5}^p-d_{2,5}d_{1,5}^p\rangle=\langle v_{14}-1,v_{13}^p-v_{12}\rangle.$$

Example 8.4. $\mathbb{F}_{p^6}$ is an a extension of both $\mathbb{F}_{p^2} = \mathbb{F}_p(b)$ and $F_{p^3} = \mathbb{F}_p(a)$. The subgroup $E := \mathrm{Span}_{\mathbb{F}_p}\{1, a, a^3, b, ba\}$ contains the subgroup $\{1, b, a, ab\}$, which is a vector space over $\mathbb{F}_{p^2}$. However $E = \mathbb{F}_{p^3} \oplus (a+b)\mathbb{F}_{p^2}$ so the partition is $(3, 2)$ rather than $(2, 2, 1)$. Note that, using Theorem 3.5, a rank 5 subgroup E is a subgroup of a dilation of $\mathbb{F}_{p^6}$ if and only if $d_{1,5}(E) \neq 0$ and $v_1(E) = v_{12}(E)^p = v_{13}(E)^{p^2} = v_{14}(E)^{p^3}$.

Using Theorem 6.2, if E is in the variety determined by

$$\langle d_{1,4}^{p^2} d_{2,4}^p d_{3,4} - d_{4,4}^p d_{1,4}^{p^2+1} - d_{3,4}^{p^2+1} \rangle \cap \mathbb{F}[d_{1,5}, \ldots, d_{5,5}]$$

then E contains a dilation of $\mathbb{F}_{p^2}$. For convenience, denote $r_1 := d_{5,5}^p - d_{4,5} d_{1,5}^p$ and $r_2 := d_{3,5}^p - d_{2,5} d_{1,5}^p$.

Conjecture 8.5.

$$\langle d_{1,4}^{p^2} d_{2,4}^p d_{3,4} - d_{4,4}^p d_{1,4}^{p^2+1} - d_{3,4}^{p^2+1} \rangle \cap \mathbb{F}[d_{1,5}, \ldots, d_{5,5}] = \langle r_1^{p^2+1} - r_2^{p^2} (d_{5,5}^p d_{2,5} - d_{4,5} d_{3,5}^p) \rangle.$$

Conjecture 8.5 has been verified for primes less than or equal to 23.

Thus we expect a subgroup E with partition $(2, 1, 1, 1)$ to satisfy

$$\left(r_1^{p^2+1} - r_2^{p^2} \left(d_{5,5}^p d_{2,5} - d_{4,5} d_{3,5}^p \right) \right)(E) = 0$$

with either $r_1(E) \neq 0$ or $r_2(E) \neq 0$.

References

[1] W. Bosma, J. J. Cannon and C. Playoust, *The Magma algebra system I: the user language*, J. Symbolic Comput. **24** (1997), 235–265.

[2] H.E.A. Campbell, I.P. Hughes, R.J. Shank and D.L. Wehlau, *Bases for rings of coinvariants*, Transformation Groups **4** (1996) 307–336.

[3] H.E.A. Campbell, R.J. Shank and D.L. Wehlau, *Rings of invariants for modular representations of elementary abelian p-groups*, Transformation Groups **18** (2013) 1–22.

[4] H. Derksen and G. Kemper, *Computational Invariant Theory*, Encyclopaedia of Math. Sci. **130**, Springer, 2002.

[5] L. E. Dickson, *An invariantive investigation of irreducible binary modular forms*, Trans. Amer. Math. Soc. **12** (1911), 1–18.

[6] C.W. Wilkerson, *A primer on the Dickson invariants*, Proceedings of the Northwestern Homotopy Theory Conference (Evanston, Ill., 1982), 421–434, Contemp. Math., **19**, Amer. Math. Soc., Providence, RI, 1983.

Department of Mathematics,
University of New Brunswick, Fredricton, NB, E3B 5A3, Canada

E-mail address: heac@unb.ca

Department of Mathematics,
University of New Brunswick, Fredricton, NB, E3B 5A3, Canada

E-mail address: jchuai@unb.ca

School of Mathematics, Statistics & Actuarial Science,
University of Kent, Canterbury, CT2 7NF, UK

E-mail address: R.J.Shank@kent.ac.uk

Department of Mathematics and Computer Science,
Royal Military College, Kingston, ON, K7K 7B4, Canada

E-mail address: wehlau@rmc.ca